David J. Pinskey et al. U.S. Serial No.:09/053,871 Filed: April 1, 1998 Page 2

--DETOXI-GEL™-

In the claims:

Please amend claim 29 as follows:

--29. (amended) A method of inhibiting clot formation in a subject which comprises adding to blood an amount of an inactive recombinant mutein in an amount to inhibit clot formation in the effective significantly subject but which does not interfere with hemostasis when the blood is administered \to a patient_

> inactive recombinant mutein wherein the comprises:

- (a) a proteolytically inactive recombinant mutein of Factor IX, or
- (b) a proteolyttically inactive recombinant mutein of Factor IXa

and wherein the recombinant mutein comprises a substitution, deletion, or addition of one or more amino acids to wild-type Factor IX or wildtype Factor IXa resulting in reduced ability to convert Factor X to Factor Xa.--

--38. (new)

The method of claim 29, wherein the recombinant mutein of Factor IX comprises an amino acid substitution for one or more of His221, Asp269 or Ser365 of wild-type Factor IX.--

34. (new)

The method of claim 33, wherein the substitution for Ser365 of wild-type Factor IX is alanine.--

David J. Pinskey et al. U.S. Serial No.:09/053,871

Filed: April 1, 1998 Page 3 ----

-35. (new)

The method \of claim 29, wherein the recombinant mutein of Factor IX comprises a deletion of one or more amino acids of wild-type Factor IX. -- :

-36. (new)

The method of claim 29, wherein the recombinant mutein of Factor IX consists essentially of amino acids 1-47 of Factor IX, and further comprises at least one\amino acid substitution from wild-type Factor IX sequence. --

(new)

The method of claim 29, wherein the substitution, deletion or addition occurs at an active site within Factor IX or Factor IXa, rendering the mutein function Ply inactive. --

. (new)

The method of claim 29, wherein the recombinant mutein of Factor IXa comprises a deletion of one or more amino acids of wild-type Factor IXa.--

8. (new)

The method of claim 2%, wherein the recombinant mutein of Factor IXa comprises an amino acid substitution for one or more of His41, Asp89 or Ser185 in the heavy chain of wild-type Factor IXa.--

3/9. (new)

The method of claim 28, wherein the substitution for Ser185 of wild-type Factor IXa is alanine.--

REMARKS

Claims 1-32 were pending. The Examiner withdrew claims 1-28 and non-elected 31 - 32from consideration as being drawn to inventions.